

DRUG FACTS

ACTIVE INGREDIENT (IN EACH 15 ML TABLESPOON): BISMUTH SUBSALICYLATE 262 MG. PURPOSE: UPSET STOMACH RELIEVER AND ANTIDIARRHEAL

USES

RELIEVES: • HEARTBURN • INDIGESTION • UPSET STOMACH • NAUSEA • DIARRHEA. (ALSO RELIEVES TRAVELER'S DIARRHEA AND GAS, BELCHING, AND FULLNESS DUE TO OVERINDULGENCE WITH FOOD AND DRINK.)

WARNINGS

REYE'S SYNDROME: CHILDREN AND TEENAGERS WHO HAVE OR ARE RECOVERING FROM CHICKEN POX OR FLU-LIKE SYMPTOMS SHOULD NOT USE THIS PRODUCT. WHEN USING THIS PRODUCT, IF CHANGES IN BEHAVIOR WITH NAUSEA OR VOMITING OCCUR, CONSULT A DOCTOR BECAUSE THESE SYMPTOMS COULD BE AN EARLY SIGN OF REYE'S SYNDROME, A RARE BUT SERIOUS ILLNESS. **ALLERGY ALERT: CONTAINS SALICYLATE.** DO NOT TAKE IF YOU ARE: • ALLERGIC TO SALICYLATES (INCLUDING ASPIRIN) • TAKING OTHER SALICYLATE PRODUCTS. **DO NOT USE IF YOU HAVE:** • AN ULCER • A BLEEDING PROBLEM • BLOODY OR BLACK STOOL. **ASK A DOCTOR BEFORE USE IF YOU HAVE:** • FEVER • MUCUS IN THE STOOL. **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE TAKING ANY DRUG FOR:** • ANTICOAGULATION (THINNING THE BLOOD) • DIABETES • GOUT • ARTHRITIS. **WHEN USING THIS PRODUCT** A TEMPORARY, BUT HARMLESS, DARKENING OF THE STOOL AND/OR TONGUE MAY OCCUR. **STOP USE AND ASK A DOCTOR IF:** • SYMPTOMS GET WORSE • RINGING IN THE EARS OR LOSS OF HEARING OCCURS • DIARRHEA LASTS MORE THAN 2 DAYS. **IF PREGNANT OR BREAST-FEEDING,** ASK A HEALTH PROFESSIONAL BEFORE USE. **KEEP OUT OF REACH OF CHILDREN.** IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.



DI 9FFT731

R731

DRUG FACTS

ACTIVE INGREDIENT (IN EACH 15 ML TABLESPOON): BISMUTH SUBSALICYLATE 262 MG. PURPOSE: UPSET STOMACH RELIEVER AND ANTIDIARRHEAL

USES

RELIEVES: • HEARTBURN • INDIGESTION • UPSET STOMACH • NAUSEA • DIARRHEA. (ALSO RELIEVES TRAVELER'S DIARRHEA AND GAS, BELCHING, AND FULLNESS DUE TO OVERINDULGENCE WITH FOOD AND DRINK.)

WARNINGS

REYE'S SYNDROME: CHILDREN AND TEENAGERS WHO HAVE OR ARE RECOVERING FROM CHICKEN POX OR FLU-LIKE SYMPTOMS SHOULD NOT USE THIS PRODUCT. WHEN USING THIS PRODUCT, IF CHANGES IN BEHAVIOR WITH NAUSEA OR VOMITING OCCUR, CONSULT A DOCTOR BECAUSE THESE SYMPTOMS COULD BE AN EARLY SIGN OF REYE'S SYNDROME, A RARE BUT SERIOUS ILLNESS. **ALLERGY ALERT: CONTAINS SALICYLATE.** DO NOT TAKE IF YOU ARE: • ALLERGIC TO SALICYLATES (INCLUDING ASPIRIN) • TAKING OTHER SALICYLATE PRODUCTS. **DO NOT USE IF YOU HAVE:** • AN ULCER • A BLEEDING PROBLEM • BLOODY OR BLACK STOOL. **ASK A DOCTOR BEFORE USE IF YOU HAVE:** • FEVER • MUCUS IN THE STOOL. **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE TAKING ANY DRUG FOR:** • ANTICOAGULATION (THINNING THE BLOOD) • DIABETES • GOUT • ARTHRITIS. **WHEN USING THIS PRODUCT** A TEMPORARY, BUT HARMLESS, DARKENING OF THE STOOL AND/OR TONGUE MAY OCCUR. **STOP USE AND ASK A DOCTOR IF:** • SYMPTOMS GET WORSE • RINGING IN THE EARS OR LOSS OF HEARING OCCURS • DIARRHEA LASTS MORE THAN 2 DAYS. **IF PREGNANT OR BREAST-FEEDING,** ASK A HEALTH PROFESSIONAL BEFORE USE. **KEEP OUT OF REACH OF CHILDREN.** IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.



DI 9FFT731

R731



DHARMA INITIATIVE BISMUTH SUBSALICYLATE

20100112CLF

NET WT. 8 FL OZ (236 mL)



DHARMA INITIATIVE BISMUTH SUBSALICYLATE

20100112CLF

NET WT. 8 FL OZ (236 mL)

DIRECTIONS

SHAKE WELL BEFORE USE • DRINK PLENTY OF CLEAR FLUIDS TO HELP PREVENT DEHYDRATION CAUSED BY DIARRHEA. • FOR ACCURATE DOSING, USE DOSE CUP. **ADULTS AND CHILDREN 12 YEARS AND OVER:** • 1 DOSE (2 TBSP OR 30 ML) EVERY 1/2 TO 1 HOUR AS NEEDED • DO NOT EXCEED 8 DOSES (16 TBSP OR 240 ML) IN 24 HOURS. • USE UNTIL DIARRHEA STOPS, BUT NOT MORE THAN 2 DAYS. **CHILDREN UNDER 12 YEARS:** ASK A DOCTOR.

OTHER INFORMATION

EACH TBSP CONTAINS: SODIUM 6 MG. • LOW SODIUM. • SUGAR FREE. • SALICYLATE 130 MG. • PROTECT FROM FREEZING • AVOID EXCESSIVE HEAT (MORE THAN 104° F OR 40° C). • TAMPER EVIDENT: DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

INACTIVE INGREDIENTS:

BENZOIC ACID, FLAVOR, MAGNESIUM ALUMINUM SILICATE, METHYLCELLULOSE, RED 22, RED 28, SACCHARIN SODIUM, SALICYLIC ACID, SODIUM SALICYLATE, SORBIC ACID, WATER



DIRECTIONS

SHAKE WELL BEFORE USE • DRINK PLENTY OF CLEAR FLUIDS TO HELP PREVENT DEHYDRATION CAUSED BY DIARRHEA. • FOR ACCURATE DOSING, USE DOSE CUP. **ADULTS AND CHILDREN 12 YEARS AND OVER:** • 1 DOSE (2 TBSP OR 30 ML) EVERY 1/2 TO 1 HOUR AS NEEDED • DO NOT EXCEED 8 DOSES (16 TBSP OR 240 ML) IN 24 HOURS. • USE UNTIL DIARRHEA STOPS, BUT NOT MORE THAN 2 DAYS. **CHILDREN UNDER 12 YEARS:** ASK A DOCTOR.

OTHER INFORMATION

EACH TBSP CONTAINS: SODIUM 6 MG. • LOW SODIUM. • SUGAR FREE. • SALICYLATE 130 MG. • PROTECT FROM FREEZING • AVOID EXCESSIVE HEAT (MORE THAN 104° F OR 40° C). • TAMPER EVIDENT: DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

INACTIVE INGREDIENTS:

BENZOIC ACID, FLAVOR, MAGNESIUM ALUMINUM SILICATE, METHYLCELLULOSE, RED 22, RED 28, SACCHARIN SODIUM, SALICYLIC ACID, SODIUM SALICYLATE, SORBIC ACID, WATER

